


View this article online at: patient.info/womens-health/polycystic-ovary-syndrome-leaflet

Polycystic ovary syndrome (PCOS)

Polycystic ovary syndrome (PCOS) is a condition that can cause symptoms of weight gain, excess hair growth, problems with periods and can impact infertility.



How common is PCOS?

PCOS is common. It is difficult to know exactly how common, as figures vary depending on the definitions used and the countries studied.

Research studies of women who had an ultrasound scan of their ovaries found that up to a third of young women have polycystic ovaries (ie ovaries with many small cysts). However, many of these women were healthy, ovulated normally and did not have high levels of male chemicals (hormones).

It is thought that around 1 in 10 women have PCOS (at least two of: polycystic ovaries, raised male hormone levels, reduced ovulation). However, the true figure may be higher because some women with mild symptoms do not seek any assessment or treatment.

[See the separate feature 8 common myths about PCOS for more details.](#)

"It is important that healthcare professionals are aware of the full range of signs of symptoms of PCOS and ensure that appropriate attention is given both to their management and their impact on an individual's mental health."

Source: Professor Adam Balen

PCOS symptoms

Symptoms typically begin in the late teens or early 20s. Not all symptoms occur in all women with PCOS. For example, some women with PCOS have some excess hair growth but have normal periods and fertility.

Symptoms of PCOS can vary from mild to severe. For example, mild unwanted hair is normal, and it can be difficult to say when it becomes abnormal in women with mild PCOS. At the other extreme, women with severe PCOS can have marked hair growth, infertility and obesity.

Symptoms may also change over the years. For example, acne may become less of a problem in middle age but hair growth may become more noticeable.

- [Period problems](#) occur in about 7 in 10 women with PCOS. You may have irregular or light periods, or no periods at all.
- [Fertility problems](#) - you need to ovulate to become pregnant. You may not ovulate each month. Some women with PCOS do not ovulate at all. PCOS is one of the most common causes of not being able to get pregnant (infertility).
- **Excess hair growth (hirsutism)** occurs in more than half of women with PCOS. It is mainly on the face, lower tummy (abdomen) and chest. In other words, it tends to be male-pattern hair. This does not happen to all women with PCOS.
- [Acne](#) may persist beyond the normal teenage years.
- [Thinning of scalp hair](#) (similar to [male pattern baldness](#)) occurs in some cases.

- **Weight gain** – women with PCOS are more at risk of becoming [overweight or obese](#).
- [Depression](#) or poor self-esteem may develop as a result of the other symptoms.

What causes PCOS?

The exact cause is not totally clear. Several factors probably play a part. These include the following:

Insulin resistance

Women with PCOS have insulin resistance. This means that cells in the body are resistant to the effect of a normal level of insulin. More insulin is then produced to keep the blood sugar normal.

This raised level of insulin causes the ovaries to make too much testosterone. A high level of insulin and testosterone causes problems with ovulation – hence, period problems and reduced fertility. It is this increased testosterone level in the blood that causes excess hair growth on the body and thinning of the scalp hair. Increased insulin also contributes towards weight gain.

Luteinising hormone (LH)

This hormone is made [in the pituitary gland, which is located in the base of the brain](#). It stimulates the ovaries to ovulate and works alongside insulin to promote testosterone production.

A high level of LH is found in about 4 in 10 women with PCOS. A high LH level combined with a high insulin level means that the ovaries are likely to produce too much testosterone.

Hereditary factors

PCOS is not usually inherited from parents but it may run in some families. There seems to be a hereditary (genetic) factor involved in some cases but this is not yet understood.

Weight

Being [overweight or obese](#) is not the underlying cause of PCOS. However, if you are overweight or obese, excess fat can make insulin resistance worse. This may then cause the level of insulin to rise even further.

High levels of insulin can contribute to further weight gain producing a 'vicious cycle'. Losing weight, although difficult, can help break this cycle.

Are any tests needed to diagnose PCOS?

Tests may be advised to clarify the diagnosis and to rule out other hormone conditions.

- **Blood tests** may be taken to measure certain chemicals (hormones). For example, a test to measure the male hormone testosterone and luteinising hormone (LH) which tend to be high in women with PCOS.
- **An ultrasound scan of the ovaries.** An ultrasound scan is a painless test that uses sound waves to create images of structures in the body. The scan can detect the typical appearance of PCOS with the many small cysts (follicles) in slightly enlarged ovaries.

The condition is diagnosed when a person has at least two of the following :

- At least 12 tiny cysts (follicles) develop in your ovaries. (Polycystic means many cysts.)
- The balance of hormones that you make in the ovaries is altered. In particular, your ovaries make more than normal of the male hormone testosterone.
- You do not ovulate each month. Some women do not ovulate at all. In PCOS, although the ovaries usually have many follicles, they do not develop fully and so ovulation often does not occur. If you do not ovulate then you may not have a period.

It is possible to have ovaries that are polycystic without having the typical symptoms that are in the syndrome. It is also possible to have PCOS without having multiple cysts in the ovary.

Screening for diabetes or pre-diabetes

Also, you may be advised to have an annual screening test for diabetes or **impaired glucose tolerance (pre-diabetes)**. A regular check for other cardiovascular risk factors such as blood pressure and blood cholesterol, may be advised to detect any abnormalities as early as possible.

Exactly when and how often the checks are done depends on your age, your weight and other factors. After the age of 40, these tests are usually recommended every three years.

PCOS treatment

There is no cure for PCOS. However, symptoms can be treated and your health risks can be reduced.

You should aim to lose weight if you are overweight

[Losing weight helps to reduce the high insulin level that occurs in PCOS](#). This has a knock-on effect of reducing the male chemical (hormone) called testosterone. This then improves the chance of you ovulating, which improves any period problems and fertility, and may also help to reduce hair growth and acne. The increased risks of long-term problems such as diabetes, high blood pressure, etc, are also reduced.

Losing weight can be difficult. A combination of eating less and [exercising more](#) is best. Advice from a dietician, and help and support from a practice nurse, may increase your chance of losing weight. Even a moderate amount of weight loss can help.

The best foods for someone with PCOS to eat are likely to be those which are slowly absorbed keeping blood sugar levels steady. These are said to have a low glycaemic index (low GI).

This means avoiding white bread, pasta and rice, and choosing wholemeal alternatives. Potatoes and sugary foods and drinks are also best avoided. Most fruit, vegetables, pulses and wholegrain foods are both healthy and have a low GI.

Occasionally tablets (such as [orlistat](#)) or [operations for weight loss](#) may be considered.

Treating hair growth

Hair growth is due to the increased level of the hormone testosterone. Unwanted hair can be removed by:

- Shaving.
- Waxing.

- Hair-removing creams.
- Electrolysis.
- Laser treatments.

These need repeating every now and then, although electrolysis and laser treatments may be more long-lasting (but are expensive and are often not available on the NHS).

There are also some medicines which may be helpful. A cream called [eflornithine](#) may be prescribed for removing unwanted facial hair. It works by counteracting a chemical (an enzyme) involved in making hair in the skin. Some research trials suggest that it can reduce unwanted hair growth, although this effect quickly wears off after stopping treatment.

Medicines taken by mouth can also treat hair growth. They work by reducing the amount of testosterone that you make, or by blocking its effect. Medicines used include:

- **Cyproterone acetate** – an anti-testosterone medicine. This is commonly combined with oestrogen as a special oral contraceptive pill called [Dianette®](#). Dianette® is commonly prescribed to regulate periods, to help reduce hair growth, to reduce acne and as a good contraceptive.
- The [combined oral contraceptive \(COC\)](#) (a combination of ethinylestradiol and drospirenone) has been shown to help if Dianette® is not suitable.
- **Other anti-testosterone medicines** are sometimes advised by a specialist if the above treatments do not help.

Medicines taken by mouth to treat hair growth take 3–9 months to work fully. You need then to carry on taking them, otherwise hair growth will come back (recur). Removing hair by the methods above (shaving, etc) may be advised whilst waiting for a medicine to work.

Treating acne

The treatments used for acne in women with PCOS are no different to the usual treatments for acne. The combined oral contraceptive pills, especially Dianette®, often help to improve acne. See the separate leaflets called [Acne](#) and [Acne Treatments](#) which covers topical treatments and antibiotic tablet treatment for acne.

Treating period problems

Some women who have no periods, or have infrequent periods, do not want any treatment for this. However, your risk of developing [endometrial cancer \(cancer of the womb/uterus\)](#) may be increased if you have no periods for a long time. Regular periods will prevent this possible increased risk to the uterus.

Therefore, some women with PCOS are advised to take the [contraceptive pill](#), as it causes regular withdrawal bleeds similar to periods. If this is not suitable, another option is to take [a progestogen hormone](#), such as medroxyprogesterone for several days every few months.

This will cause a monthly bleed like a period. Sometimes, an [intrauterine system \(IUS\)](#), which releases small amounts of progestogen into the womb, preventing a build-up of the lining, can be used. If none of these methods is suitable, your doctor may advise a regular ultrasound scan of your uterus to detect any problems early.

Fertility issues

Although fertility is often reduced, you still need [contraception](#) if you want to be sure of not getting pregnant. The chance of becoming pregnant depends on how often you ovulate. Some women with PCOS ovulate now and then, others not at all.

If you do not ovulate but want to become pregnant then [fertility treatments](#) may be recommended by a specialist and have a good chance of success. Tablets such as [clomifene](#) can cause you to ovulate.

But remember, you are much less likely to become pregnant if you are obese. If you are obese or overweight then losing weight is advised in addition to other fertility treatments.

Metformin and other insulin-sensitising medicines

Metformin is a medicine that is commonly used to treat people with type 2 diabetes. It makes the body's cells more sensitive to insulin. This may result in a decrease in the blood level of insulin which may help to counteract the underlying cause of PCOS – see above.

For certain people with PCOS, a specialist may advise that metformin be taken. However, further research is needed to confirm the role of these medicines in the treatment of PCOS.

Possible long-term problems of polycystic ovary syndrome

If you have PCOS, over time you have an increased risk of:

- Developing **type 2 diabetes**.
- Developing **diabetes in pregnancy**.
- A **high cholesterol level**.
- **High blood pressure**.
- Being **overweight**, particularly around the tummy.

These problems in turn *may* also increase your risk of having a **stroke** and **heart disease** in later life. These increased health risks are due to the long-term insulin resistance.

A sleeping problem called **sleep apnoea** is also more common than average in women with PCOS.

Other possible problems in pregnancy include more chance of having **premature babies** or having **high blood pressure in pregnancy (pre-eclampsia)**. There may be twice the risk of developing diabetes in pregnancy if you have PCOS so you would be checked for this regularly.

If you have no periods, or very infrequent periods, you *may* have a higher-than-average risk of developing **cancer of the womb (uterus)**. However, the evidence for this is not conclusive and, if there is a risk, it is probably small and can be prevented.

Preventing long-term problems

A healthy lifestyle is important to help prevent the conditions listed above in 'Possible long-term problems of polycystic ovary syndrome (PCOS)'. For example, you should:

- Eat a healthy diet.
- Exercise regularly.
- Lose weight if you are overweight or obese.
- Not smoke.

Healthy lifestyle advice applies to everyone, whether they have PCOS or not. However, it is particularly important for women with PCOS, as they may have extra risk factors for health problems in later life. These risks are much reduced if you are not overweight and do not smoke.

Further reading

- [Fertility – Assessment and treatment for people with fertility problems](#); NICE Guidance (February 2013, updated September 2017)
- [Long-term Consequences of Polycystic Ovary Syndrome](#); Royal College of Obstetricians and Gynaecologists (November 2014)
- [Hirsutism](#); NICE CKS, July 2020 (UK access only)
- [Lim SS, Hutchison SK, Van Ryswyk E, et al](#); Lifestyle changes in women with polycystic ovary syndrome. *Cochrane Database Syst Rev.* 2019 Mar 28;3:CD007506. doi: 10.1002/14651858.CD007506.pub4.
- [Fraison E, Kostova E, Moran LJ, et al](#); Metformin versus the combined oral contraceptive pill for hirsutism, acne, and menstrual pattern in polycystic ovary syndrome. *Cochrane Database Syst Rev.* 2020 Aug 13;8:CD005552. doi: 10.1002/14651858.CD005552.pub3.
- [Graff SK, Mario FM, Ziegelmann P, et al](#); Effects of orlistat vs. metformin on weight loss-related clinical variables in women with PCOS: systematic review and meta-analysis. *Int J Clin Pract.* 2016 Jun;70(6):450–61. doi: 10.1111/ijcp.12787. Epub 2016 May 26.
- [Morley LC, Tang T, Yasmin E, et al](#); Insulin-sensitising drugs (metformin, rosiglitazone, pioglitazone, D-chiro-inositol) for women with polycystic ovary syndrome, oligo amenorrhoea and subfertility. *Cochrane Database Syst Rev.* 2017 Nov 29;11:CD003053. doi: 10.1002/14651858.CD003053.pub6.
- [Polycystic ovary syndrome](#); NICE CKS, April 2024 (UK access only)

Disclaimer: This article is for information only and should not be used for the diagnosis or treatment of medical conditions. Egton Medical Information Systems Limited has used all reasonable care in compiling the information but makes no warranty as to its accuracy. Consult a doctor or other healthcare professional for diagnosis and treatment of medical conditions. For details see our [conditions](#).

Authored by:	Peer Reviewed by: Dr Krishna Vakharia, MRCGP	
Originally Published: 19/11/2023	Next review date: 19/05/2023	Document ID: doc_4585

View this article online at: patient.info/womens-health/polycystic-ovary-syndrome-leaflet

Discuss Polycystic ovary syndrome (PCOS) and find more trusted resources at [Patient](#).



To find out more visit www.patientaccess.com
or download the app



Follow us

